

Cirsium semzinanicum (Asteraceae), a new species from Hakkâri, Turkey

Mehmet Firat¹

¹ *Yüzüncü Yıl University, Faculty of Education, Department of Biology, TR-65080, Van, Turkey*

Corresponding author: Mehmet Firat (kuyucak65@yahoo.com)

Academic editor: Peter de Lange | Received 5 April 2016 | Accepted 26 July 2016 | Published 5 August 2016

Citation: Firat M (2016) *Cirsium semzinanicum* (Asteraceae), a new species from Hakkâri, Turkey. PhytoKeys 68: 65–72. doi: 10.3897/phytokeys.68.8745

Abstract

Cirsium semzinanicum **sp. nov.** (Asteraceae) is described as a new species from Hakkâri, Turkey. The new species is a part of the sect. *Epitrachys* (Cardueae) and similar to *Cirsium karduchorum*, from which it differs in morphological characters such as leaves, involucre, phyllaries, corolla, achenes and pollen morphology. Geographical distribution, habitat and IUCN conservation status of this species are given.

Keywords

Asteraceae, *Epitrachys*, *Cirsium*, Hakkari, Turkey, taxonomy

Introduction

The genus *Cirsium* Mill. is one of the largest genera of Asteraceae, and it comprises more than 250 perennial, biennial, or rarely annual spiny species distributed in the northern hemisphere in Europe; North Africa; East, Central, and Southwest Asia; and North and Central America (Charadze 1963, Davis and Parris 1975, Petrak 1979, Kadereit and Jeffrey 2007).

The last revision on *Cirsium* species growing in Turkey was carried out by Davis and Parris (1975) for the Flora of Turkey. In that study, 52 species (65 taxa) were given under 3 sections [*Cirsium*, *Epitrachys* DC., and *Cephalonoplos* (Neck.) DC.]. Additional taxonomic treatments have dealt with the distribution of the genus in supplements and 5 new species (6 taxa) were given (Davis et al. 1988; Güner et al. 2000).

Finally, the Turkish members of the genus were established as 66 species (78 taxa) according to Yıldız (2012) and Yıldız et al. (2013).

Between 2011 and 2015 new populations of *Cirsium* sp. similar to *C. karduchorum* were discovered at 3 sites in south-east Turkey. Here the morphological and micro-morphological characters of these new populations are presented and their taxonomic treatment is proposed.

Materials and methods

The specimens were collected by the author in July 2011 and August 2015 during the field trips to Şemdinli (Fig. 1) in Hakkâri, in the southeast of Turkey. In total 10 herbarium specimens of the presumed new species were collected from three adjacent localities and deposited in the herbaria of VANE, ANK and E. A formal name is provided accompanied by a detailed description and illustration.

At first glance, on the basis of the characters of leaves, involucre, phyllaries, corolla and achenes, seemed to be similar to *C. karduchorum*. The newly collected specimens were therefore cross-checked with the keys provided by de Candolle (1838), Boissier (1875), Petrak (1910), Davis and Parris (1975), Davis et al. (1988) and Güner et al. (2000). They were compared with the related specimens stored in VANE, GAZI, ANK and HUB herbaria.




Figure 1. Distribution map of *Cirsium semzinanicum* sp. nov. (▲) and also closely related species *C. karduchorum* (■) in Turkey.

Images of the collected material were taken with a Sony DSCR1, digital camera. The SEM micrographs were taken with a ZEISS supra55. The terminologies for pollen morphology were used in accordance with Faegri and Iversen (1989) and Punt et al. (1994). Geographical positions were identified using a Magellan explorer 710 GPS. According to the grid system (Davis 1965), especially the new species which are present in Hakkâri province falls within the C10 square (Davis 1965). The conservation status of the new species was assessed according to IUCN criteria (IUCN 2014).

Taxonomy treatment

Cirsium semzinanicum Firat, sp. nov.

urn:lsid:ipni.org:names:77156986-1

Figs 2–3

Type. Turkey. C10 Hakkâri: Şemdinli, Bêgoz village, Kaduna region, rocky slopes, eroded slopes, 37°17'15"N, 44°25'25"E, 1717 m, 26 July 2011, *M. Firat* 27257 (holotype VANF!, isotypes ANKA, E, and Hb. M. Firat).

Diagnosis. *Cirsium semzinanicum* clearly differs from *C. karduchorum* Petr. in its stems 100–150 cm tall (vs. 50–100 cm tall, robust), basal leaves 15–25 × 9–14 cm “excluding 10–18 petiole”, green, (vs. 30–40 × 15–20 cm “excluding 15–20 cm petiole”, bluish-green), involucre 15–25 × 20–30 mm ovoid to subglobose (vs. 25–30 × (30–)40–50 mm depressed subglobose to broadly obovoid), phyllaries 9–12-seriate ± erecto-reflexed (vs. 11–14-seriate reflexed or recurved), achen brownish 7–9 × 3–4 mm (vs. bright grayish brown 6–7.5 × 3 mm).

Description. Perennial, few stemmed from base, erect, 100–150 cm tall, unwinged but longitudinally striate, glabrous–glaucous, with few branches above, sterile shoots at the base. Basal leaves long-petiolate, petioles 10–18 cm long, winged, with large auricles; lamina broadly elliptic in outline, 15–25 × 9–14 cm, green, very sparsely spinulose–strigose above with 0.5–0.9 mm long, adpressed setae, ca 1–3 per 2 mm square, otherwise densely arachnoid; lower surface very sparsely arachnoid, 3/4 pinnatifid with 4–5-paired, 5–8 × 2–5 lateral and terminal lobes with stout 7–12 mm apical spine. Middle cauline leaves similar but smaller to basal leaves, petioles 1–6 cm long, lamina broadly elliptic in outline 4–8 × 2.5–4 cm long, lateral and terminal lobes 3/4 pinnatifid with 3–5 lobes, spinulose–strigose above with 0.7–1.2 mm long, adpressed setae, ca 4–6 per 2 mm square, otherwise densely arachnoid with stout 6–9 mm apical spine. Upper cauline leaves similar middle cauline leaves but smaller and sessile, shortly auriculate. Uppermost (involucral) leaves 2–3, 1–3 cm long, linear–lanceolate with sinuate–dentate margin, shorter than involucre, sessile. Capitula erect, 1(–2) on each branch, compound corymbose with 15–30–capitula, 26–35 mm × 2–30 mm, sessile or peduncles 1–2 cm long; involucre 15–25 × 20–30 mm, ovoid to subglobose; phyllaries 9–12-seriate, adpressed, glabrous, greenish–yellowish, ± erecto-reflexed, the outer 8–10 × 2–3 mm, with reflexed, ovate-oblong, 3–5 × 0.3–0.4 mm long apical spine;




Figure 2. *Cirsium semzinanicum* sp. nov.: **a** habits **b** basal leaf **c** upper leaf surface **d** capitulum. *Cirsium karduchorum*: **e** habit **f** basal leaf **g** upper leaf surface **h** capitulum.

the median $10\text{--}14 \times 2\text{--}3$ mm long, with \pm reflexed, oblong-lanceolate, $4\text{--}6 \times 0.2\text{--}0.3$ mm long apical spine; the inner $16\text{--}21 \times 1\text{--}2$ mm, linear-lanceolate, $5\text{--}7 \times 0.1\text{--}0.2$ mm long apical spine. Corolla pinkish-whitish, 26–32 mm long, lobed to $1/4\text{--}1/5$.




Figure 3. *Cirsium semzinanicum* sp. nov.: **a** outer phyllary **b** median phyllary **c** inner phyllary **d** achene. *Cirsium karduchorum*: **e** outer phyllary **f** median phyllary **g** inner phyllary **h** achene.

Style 15–19 mm long, exerted, shortly bilobed; filaments 5–7 mm long, densely hairy; Achenes brownish, 7–9 × 3–4 mm, oblong, asymmetric, slightly compressed. Pappus 16–18 mm long, plumose, dirty white, or light brown.

Micromorphology. Pollen grains are tricolporate, pollen shapes P/E: spheroidal, amb circular, polar axis $36.72\ \mu\text{m}$, equatorial axis $36.60\ \mu\text{m}$, exine $1.62\ \mu\text{m}$ thick, ornamentation echinate. Tectum complete structured, spines conic and pointed, 5–6 per $100\ \mu\text{m}^2$, $2.83\ \mu\text{m}$ long, base diameter $3.45\ \mu\text{m}$, intine $1.02\ \mu\text{m}$, colpi margins are straight and distinct with pointed ends, $10\ \mu\text{m}$ long (Fig. 4).

Pollen grains are tricolporate, pollen shapes P/E: spheroidal, amb circular, polar axis $36.21\ \mu\text{m}$, equatorial axis $38.25\ \mu\text{m}$, exine $1.73\ \mu\text{m}$ thick, ornamentation echinate, tectum complete structured, perforate with spines conic and pointed, 5–6 per $100\ \mu\text{m}^2$, $2.70\ \mu\text{m}$ long, base diameter $3.45\ \mu\text{m}$, intine $0.95\ \mu\text{m}$, colpi margins are straight and distinct with pointed ends, $17.5\ \mu\text{m}$ long (Fig. 5).




Figure 4. SEM Microphotograph of pollen of *Cirsium karduchorum*.



Figure 5. SEM Microphotograph of pollen of *Cirsium semzinanicum* sp. nov.

Phenology. Flowering and fruiting, from July to September.

Vernacular name. *Cirsium* spp. are called as “Kîvar” by the local people of the Hakkâri province (Firat 2013).

Etymology. The specific epithet is derived from the name of the Şemzinan (Şemdinli) province where type material was collected.

Distribution and conservation status. *Cirsium semzinanicum* is endemic to Hakkâri province in Turkey. The number of mature individuals is approximately 600 and is known from 3 locations (criteria B2ab [i.iii]). Therefore, it should be regarded as belonging to the IUCN Vulnerable (VU) threat category (IUCN 2014).

Habitat and ecology. The new species grows in Oak openings, slopes and eroded slopes at c. 1600–1900 m elevation with plants such as *Quercus libani*, *Daphne oleoides* subsp. *kurdica*, *Satureja bachtiarica*, *Dianthus orientalis*, *Echinops tournefortii*, *Astragalus* sp., *Eryngium billardierei*.

Discussion

Cirsium semzinanicum is morphologically similar to *C. karduchorum* because of having similar habitus, lesser involucre and phyllaries, corolla and achen size, but it is easily distinguished from its stems 100–150 cm tall (not 50–100 cm tall, robust), basal leaves 15–25 × 9–14 cm “exculiding 10–18 petiole”, green, (not 30–40 × 15–20 cm “exculiding 15–20 petiole”, bluish-green), involucre 15–25 × 20–30 mm ovoid to subglobose (not 25–30 × (30–)40–50 mm depressed subglobose to broadly obovoid), phyllaries 9–12-seriate ± erecto-reflexed (not 11–14-seriate reflexed or recurved), achen brownish 7–9 × 3–4 mm (not bright grayish brown 6–7.5 × 3 mm). Additional morphological differences between the new species and *C. karduchorum* are given in Table 1. Pollen morphology of both of species shows similarity. Pollen grains are isopolar, tricolporate, pollen shapes are spheroidal. Colpus length of *C. semzinanicum* is longer than *C. karduchorum*, while tectum completely structured, spines conic and pointed in *C. karduchorum*, the tectum complete structured, perforate with spines conic and pointed in *C. semzinanicum* (Figs 4 and 5).

Additional specimens examined

Cirsium semzinanicum: Turkey. C10 Hakkâri: Şemdinli, above Benavok village, Gerdi şapatan region, rocky slopes, eroded slopes, 37°09'53"N, 44°26'00"E, 1692 m, 8 August 2013, *M. Fırat* 30332 (VANF! and Hb. M. Fırat); C10 Hakkâri: Şemdinli, Berxoşe region, rocky slopes, 37°20'62"N, 44°34'64"E, 2072 m, 4 August 2014, *M. Fırat* 30958 (Hb. M. Fırat)

Cirsium karduchorum: Turkey. C9 Hakkâri: Karadağ mountain above Hakkâri, eroded slopes, 2438 m, 13 August 1954, *Davis & Polunin* 24326 (ANK!, E! and W!); Hakkâri: from Hakkâri to Berçelan plateau, eroded slopes, 2150 m, 5 September 2007, *M. Fırat & T. Dirmenci* 3579 (Hb. M. Fırat); Hakkâri: from Hakkâri to Berçelan plateau, eroded slopes, 37°36'00"N, 43°44'50"E, 2130 m, 16 August 2008, *M. Fırat, T. Dirmenci, B. Yıldız* 16932 (Hb. M. Fırat); Hakkâri: Kotranis region, eroded slopes, 37°44'12"N, 43°44'51"E, 2277 m, 16 August 2011, *M. Fırat* 27767 (Hb. M. Fırat); Hakkâri: from Hakkâri to Berçelan plateau, eroded slopes, 37°36'00"N, 43°44'50"E, 2130 m, 16 August 2013, *M. Fırat* 30439 (Hb. M. Fırat).

Acknowledgements

Thanks are due to Dr. Rıza Binzet for helping with pollen studies.

References

- Boissier E (1875) *Cirsium* Mill. In: Boissier E (Ed.) *Flora Orientalis*, Vol. 3. Geneva, 523–553.
- Candolle AP (1838) *Prodromus Systematis Naturalis Regni Vegetabilis*, Vol. 6. Treuttel et Würtz, Paris.
- Charadze AL (1963) *Cirsium* Mill. In: Bobrov EG, Cherepanov SK (Eds) *Flora of the USSR*, Vol. XXVIII. Izdatel'stvo Akademii Nauk SSSR, Moscow/Leningrad, 63–270. [Translated by Russian–Israel Program for Scientific Translations, Jerusalem (English version, 1976)]
- Davis PH, Mill RR, Tan K (1988) *Flora of Turkey and the East Aegean Islands* (Suppl. 1), Vol. 10. Edinburgh University Press, Edinburgh, 164–165.
- Davis PH, Parris BS (1975) *Cirsium* Mill. In: Davis PH (Ed.) *Flora of Turkey and the East Aegean Islands*, Vol. 5. Edinburgh University Press, Edinburgh, 370–412.
- Davis PH (1965) *Flora of Turkey and the East Aegean Islands*, Vol. 1. Edinburgh University Press, Edinburgh, 567 pp.
- Fægri K, Iversen J (1989) *Textbook of pollen analysis*. John Wiley, Sons, Chichester, 328 pp.
- Fırat M (2013) *Ferhenga Navên Riwekên Bi Kurdî / Kürtçe Bitki Adları Sözlüğü/Dictionary of Plant Names in Kurdish*. Kalkan Ofset, Ankara, 652 pp.
- Güner A, Özhatay N, Ekim T, Başer KHC (2000) *Flora of Turkey and the East Aegean Islands* (Suppl. 2), Vol. 11, Edinburgh University Press, Edinburgh.
- IUCN (2014) *IUCN Red List Categories and Criteria: Version 11*. IUCN Species Survival Commission, Gland, Switzerland and Cambridge, 87 pp.
- Kadereit JW, Jeffrey C (2007) *The Families and Genera of Vascular Plants*, Vol. 8. Flowering Plants. Eudicots: Asterales. Springer, Berlin. doi: 10.1007/978-3-540-31051-8
- Petrak F (1910) Über neue oder wenig bekannte Cirsien aus dem Orient. *Oesterreichische Botanische Zeitschrift* 60: 393–396. doi: 10.1007/BF01647979
- Petrak F (1979) *Cirsium* Mill. In: Rechinger KH (Ed.) *Flora Iranica. Compositae III-Cynareae*, Vol. 139a. Akademische Druck-u Verlagsanstalt, Graz, 231–280.
- Punt W, Blackmore S, Nilsson S, Le Thomas A (1994) *Glossary of Pollen and Spore Terminology*. LPP Foundation, Utrecht, 71 pp.
- Yıldız B (2012) *Cirsium* Mill. In: Güner A, Aslan S, Ekim T, Vural M, Babaç MT (Eds) *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, İstanbul, 141–146.
- Yıldız B, Arabacı T, Dirmenci T (2013) Twonew species of *Cirsium* (Asteraceae) and notes on allies from Turkey, *Turkish Journal of Botany* 37: 1045–1054.